Florida Citrus Advanced Technology Program

QUARTERLY & FINAL REPORTS: Control of Citrus Greening, Canker & Emerging Diseases of Citrus

Instructions Complete the fields based on your project specs. When finished, save the form to your local disk using a unique name. Then, go to **http://research.fcprac.com**, and log in with your user name and password using Researcher Login in the lower left. Find this project title and click on **Submit a Report**. Update your profile information if needed, then upload this report as directed.

2009-2010 REPORT	CATEGORY (drop-down)	TODAY'S DATE (m/d/yr)
◉ *Quarterly Report* ○ *Annual Report* ○ *Final*	Psyllid	10/15/09

WHAT IS THE "HEADLINE" FOR THIS REPORT (e.g. a one-sentence "newspaper headline" describing what you accomplished)

Insecticides interfere with the probing behavior of Diaphorina citri and is efficient in controlling

TITLE and CONTACT INFORMATION

Proposal Title CAN INSECTICIDES AND MINERAL OIL AVOID TRANSMISSION OF Candidatus Liberibacter asiaticus BY Diaphorina citri?

Principal Investigator Pedro Takao Yamamoto	*PI Last Name* Yamamoto
Email ptyamamoto@fundecitrus.com.br	*FDACS Contract Number* 236
Phone +55 16 3301-7025	*Project Duration (years)* 2 *Year of Project* 1
Organization Fundecitrus	*Total Direct Funds (current year)* US$ 15,710.00

REPORT UPDATE (650 words; provide details about your headline)

In the first experiment that was conducted to know the efficiency of systemic insecticides to control the Asian Citrus Psyllid (ACP), Diaphorina citri, and its effect on transmission of the bacteria Candidatus Liberibacter asiaticus, indicated that imidacloprid (Confidor 700 GrDA), 0.35 g AI/plant and thiamethoxam (Actara 250 WG) 0.25 g AI/plant, applied in the nursery tree bags, before planting, was efficient to control ACP until 60 days after application. The time to cause 100% of ACP mortality was between 5 to 7 days after the confinement of adults in treated plants. However, researches using electrical penetration graph (EPG) showed that in plants treated with imidacloprid and thiamethoxam, after the first feeding on phloem, the adults do not do more probing. No transmission results yet. We started the second experiment to determine if the systemic insecticides are effective until 90 days after application and its effect on transmission of the bacteria.

Using electrical penetration graphs (EPG) techniques, we are studying the probing behavior of ACP. In the preliminary results we observed that the time feeding on phloem, in plants which were applied the systemic insecticides imidacloprid, was 7.58 and 6.95 minutes to 35 and 95 days after application, respectively. When applied thiamethoxam the results were 13.28, 15.41, and 17.50 minutes, respectively to 20, 35 and 95 days after application. In the control, the average time feeding on phloem was 129.79 minutes. For both insecticides, all specimens died in two days after those times of phloem sap ingestion. After first probing and ingestion of phloem sap, the insect stopped feeding and no more probing was observed.

In foliar application, were tested imidacloprid (Provado 200 SC) and lambda-cyhalothrin (Karate 50 SC). Until 14 days after application, adults of ACP probe the citrus trees, but did not reach the phloem and died in 24 hours. In the test plant, approximately 80% of the ACP adults reached and sucked on phloem for 2 hours, in average, in a 5-hour record test.